

FEC : ETAT DES LIEUX SUR 8 ETABLISSEMENTS BRETONS 2008

	DESFERAZOXANE			EPIRUBICINE			CYCLOPHOSPHAMIDE			5 FU			durée FEC seul	observations
	V	mg/m2	durée	V	mg/m2	durée	V	mg/m2	durée	V	mg/m2	durée	mn	
A	250ml	1000	15mn	100ml G5%	100	10mn	100ml Nacl 0,9%	500	15mn	20ml nacl 0,9%	500	2mn	27	
B	200ml	1000	20mn	seringue	100	20mn	100ml G5%	500	30mn	20ml nacl 0,9%	500	5mn	55	
C	200ml	1000	15mn	100ml G5%	100	15mn	100ml Nacl 0,9%	500	15mn	20ml nacl 0,9%	500	2mn	32	
D				seringue	100	ivl	100ml Nacl 0,9%	500	90mn	100ml Nacl 0,9%	500	90mn	92	F avant E
E				seringue	100	20mn	250ml G5%	500	60mn	250ml G5%	500	60mn	80	mélange F+C
F				100ml Nacl 0,	100	10mn	100ml G5%	500	10mn	250ml Nacl 0,9%	500	60mn	80	
G				100ml G5%	100	15mn	100ml G5%	500	15mn	100ml G5%	500	15mn	45	
H	200ml	1000	20mn	100ml G5%	100	10mn	100ml G5%	500	30mn	100ml G5%	500	30mn	80	
*				100ml G5%	100	15mn	100ml G5%	500	60mn	100ml G5%	500	15mn	90	

PACS 04/0005 bras A pas de données précises

PACS 05/0106 "IV lente dans la tubulure d'une perfusion à débit rapide de G5%"

*BONNETERRE J. et al. Indication : cancer du sein A.S.C.O. 1998, abst. 473